

Repellent activity of catmint, *Nepeta cataria*, and iridoid nepetalactone isomers against Afro-tropical mosquitoes, ixodid ticks and red poultry mites

Michael A. Birkett^{a,*}, Ahmed Hassanali^{b,c}, Solveig Hoglund^d, Jan Pettersson^d, John A. Pickett^a

^a Biological Chemistry Department, Rothamsted Research, Harpenden, Hertfordshire AL5 2JQ, UK

^b Behavioural and Chemical Ecology Department, International Centre for Insect Physiology and Ecology, P.O. Box 30772-00100, Nairobi, Kenya

^c Chemistry Department, Kenyatta University, P.O. Box 43844, Nairobi, Kenya

^d Department of Ecology, Swedish University of Agricultural Sciences, P.O. Box 7044, SE-750 07 Uppsala, Sweden

ARTICLE INFO

Article history:

Received 28 July 2010

Accepted 26 September 2010

Available online 4 November 2010

Keywords:

Nepeta cataria

Anopheles gambiae

Culex quinquefasciatus

Rhipicephalus appendiculatus

Dermanyssus gallinae

Repellent

Nepetalactones

(*E*)-(1*R*,9*S*)-Caryophyllene

ABSTRACT

The repellent activity of the essential oil of the catmint plant, *Nepeta cataria* (Lamiaceae), and the main iridoid compounds (4*aS*,7*S*,7*aR*) and (4*aS*,7*S*,7*aS*)-nepetalactone, was assessed against (i) major Afro-tropical pathogen vector mosquitoes, i.e. the malaria mosquito, *Anopheles gambiae* s.s. and the Southern house mosquito, *Culex quinquefasciatus*, using a World Health Organisation (WHO)-approved topical application bioassay (ii) the brown ear tick, *Rhipicephalus appendiculatus*, using a climbing repellency assay, and (iii) the red poultry mite, *Dermanyssus gallinae*, using field trapping experiments. Gas chromatography (GC) and coupled GC–mass spectrometry (GC–MS) analysis of two *N. cataria* chemotypes (A and B) used in the repellency assays showed that (4*aS*,7*S*,7*aR*) and (4*aS*,7*S*,7*aS*)-nepetalactone were present in different proportions, with one of the oils (from chemotype A) being dominated by the (4*aS*,7*S*,7*aR*) isomer (91.95% by GC), and the other oil (from chemotype B) containing the two (4*aS*,7*S*,7*aR*) and (4*aS*,7*S*,7*aS*) isomers in 16.98% and 69.83% (by GC), respectively. The sesquiterpene hydrocarbon (*E*)-(1*R*,9*S*)-caryophyllene was identified as the only other major component in the oils (8.05% and 13.19% by GC, respectively). Using the topical application bioassay, the oils showed high repellent activity (chemotype A $RD_{50} = 0.081 \text{ mg cm}^{-2}$ and chemotype B $RD_{50} = 0.091 \text{ mg cm}^{-2}$) for *An. gambiae* comparable with the synthetic repellent DEET ($RD_{50} = 0.12 \text{ mg cm}^{-2}$), whilst for *Cx. quinquefasciatus*, lower repellent activity was recorded (chemotype A $RD_{50} = 0.34 \text{ mg cm}^{-2}$ and chemotype B $RD_{50} = 0.074 \text{ mg cm}^{-2}$). Further repellency testing against *An. gambiae* using the purified (4*aS*,7*S*,7*aR*) and (4*aS*,7*S*,7*aS*)-nepetalactone isomers revealed overall lower repellent activity, compared to the chemotype A and B oils. Testing of binary mixtures of the (4*aS*,7*S*,7*aR*) and (4*aS*,7*S*,7*aS*) isomers across a range of ratios, but all at the same overall dose (0.1 mg), revealed not only a synergistic effect between the two, but also a surprising ratio-dependent effect, with lower activity for the pure isomers and equivalent or near-equivalent mixtures, but higher activity for non-equivalent ratios. Furthermore, a binary mixture of (4*aS*,7*S*,7*aR*) and (4*aS*,7*S*,7*aS*) isomers, in a ratio equivalent to that found in chemotype B oil, was less repellent than the oil itself, when tested at two doses equivalent to 0.1 and 0.01 mg chemotype B oil. The three-component blend including (*E*)-(1*R*,9*S*)-caryophyllene at the level found in chemotype B oil had the same activity as chemotype B oil. In a tick climbing repellency assay using *R. appendiculatus*, the oils showed high repellent activity comparable with data for other repellent essential oils (chemotype A $RD_{50} = 0.005 \text{ mg}$ and chemotype B $RD_{50} = 0.0012 \text{ mg}$). In field trapping assays with *D. gallinae*, addition of the chemotype A and B oils, and a combination of the two, to traps pre-conditioned with *D. gallinae*, all resulted in a significant reduction of *D. gallinae* trap capture. In summary, these data suggest that although the nepetalactone isomers have the potential to be used in human and livestock protection against major pathogen vectors, intact, i.e. unfractionated, *Nepeta* spp. oils offer potentially greater protection, due to the presence of both nepetalactone isomers and other components such as (*E*)-(1*R*,9*S*)-caryophyllene.

© 2010 Elsevier Ltd. All rights reserved.

1. Introduction

Vector-borne infectious diseases affecting livestock and human health contribute directly to food insecurity, poverty and mortality on a global scale (FAO, 2005; WHO, 2000). Therefore, the availability of a range of tools for control of pathogen vectors, predomi-

* Corresponding author. Tel./fax: +44 1582 763133.

E-mail address: mike.birkett@bbsrc.ac.uk (M.A. Birkett).

nantly invertebrate arthropods (mosquitoes, ticks and mites), is of the utmost importance. Arthropods rely predominantly upon olfactory perception of volatile, host-derived semiochemicals (behaviour-modifying chemicals) during host location (Logan and Birkett, 2007), and, therefore, whilst the potential for deploying broad-spectrum neurotoxins, i.e. pesticides, in vector control has been reduced in recent years, the nervous system still remains a major target for intervention, through behaviour modification. Thus, protection against arthropod biting and blood-sucking behaviour through the development of topically-applied and spatial repellents has received much attention in recent years (Debboun et al., 2006), with development being based upon three hypotheses that relate to the evolution of repellency: (i) the avoidance of strong plant cues from botanicals, (ii) the avoidance of closely related, but taxonomically different, unsuitable hosts and (iii) the avoidance of highly resistant individuals within a host species complex (Pickett et al., 2010).

A number of repellents have been identified from botanical sources, based on prior ethnobotanical use, thus exploiting the natural ability of such compounds to interfere with host location. These include citronella oil, *Eucalyptus* spp. oils containing *p*-menthane-3,8-diol and eucamadol, and gum resins from members of the Burseraceae, which have been investigated for use as repellents for human or livestock protection (Nishimura and Satoh, 1999; Peterson and Coats, 2001; Birkett et al., 2008). However, most of the active components from botanical sources are highly volatile and can be readily lost, therefore losing efficacy (Lindsay et al., 1996; Chou et al., 1997). This loss of activity has resulted in such materials being overlooked in favour of synthetic repellents such as DEET, KBR3023 and IR3535 that were developed through structure–activity relationship studies (Debboun et al., 2006). Botanically-derived repellents have been discovered which may avoid problems with loss of activity, and which may overcome the negative impact of synthetic repellents. These include isoprenoids from *Callicarpa* spp. (Verbenaceae) (Cantrell et al., 2005; Carroll et al., 2007) and the isoprenoid-derived compound isolongifolone (Zhang et al., 2009). Much attention has focused on the catmint plant, *Nepeta cataria* (Lamiaceae), and its major essential oil components, i.e. iridoid nepetalactones, of which the (4*a*S,7*S*,7*a*R) and (4*a*S,7*S*,7*a*S) isomers predominate. Repellent activity has been reported against mosquitoes (Chauhan et al., 2005; Bernier et al., 2005; Schultz et al., 2006). Hydrogenated *N. cataria* oil enriched in dihydronepetalactone isomers has been reported to be effective against mosquitoes and blackflies (Spero et al., 2008). The activity of *N. cataria* oil against *Aedes*, *Anopheles* and *Culex* spp. mosquitoes in North Africa, Australia and the Far East has been reported (Amer and Mehlhorn, 2006; Webb and Russell, 2007; Polsomboon et al., 2008). However, repellent activity against either sub-Saharan populations of Afro-tropical mosquito species or global populations of acarine pests, i.e. ixodid ticks and mites, has not been reported.

Based on existing ethnobotanical principles, the repellent activity of essential oils from several plant families, including members of the Lamiaceae family, has been assessed against sub-Saharan populations of Afro-tropical mosquito species and ixodid ticks (Omolo et al., 2004; Odalo et al., 2005; Mwangi et al., 1995; Ndungu et al., 1995; Lwande et al., 1999). Thus, the aim of this study was to assess the repellent activity of oils from two chemotypes of *N. cataria*, i.e. another member of the Lamiaceae, enriched in (4*a*S,7*S*,7*a*R) and (4*a*S,7*S*,7*a*S)-nepetalactone isomers (Birkett and Pickett, 2003) against the same target species, using a WHO-approved topical application assay (WHO, 1996) and a tick-climbing repellency bioassay (Wanzala et al., 2004). The study also investigated the repellent activity of (4*a*S,7*S*,7*a*R) and (4*a*S,7*S*,7*a*S) isomers versus another acarine pest, the red poultry mite, *Dermanyssus gallinae*, in order to determine their potential for use as general acarine repellents.

2. Results

2.1. *Nepeta cataria* oil composition

Coupled GC–MS analysis of the oils revealed the presence of (4*a*S,7*S*,7*a*R) and (4*a*S,7*S*,7*a*S)-nepetalactone isomers, respectively, with one of the oils (chemotype A) being dominated by the (4*a*S,7*S*,7*a*R) isomer (91.95% by GC), and the other oil (chemotype B) containing both (4*a*S,7*S*,7*a*R) and (4*a*S,7*S*,7*a*S) isomers (16.98% and 69.83% by GC, respectively) (Fig. 1). The sesquiterpene hydrocarbon (*E*)-(1*R*,9*S*)-caryophyllene was also identified in chemotype A and B oils (8.05% and 13.19%, respectively) (Fig. 1).

2.2. Repellency assays with essential oil samples

Using a WHO-approved topical application bioassay, the *N. cataria* oils showed high repellent activity (chemotype A $RD_{50} = 0.081 \text{ mg cm}^{-2}$ and chemotype B $RD_{50} = 0.091 \text{ mg cm}^{-2}$) for *Anopheles gambiae* comparable with the synthetic repellent DEET ($RD_{50} = 0.12 \text{ mg cm}^{-2}$) (Fig. 2), whilst for *Culex quinquefasciatus*, lower repellent activity was recorded (chemotype A $RD_{50} = 0.34 \text{ mg cm}^{-2}$ and chemotype B $RD_{50} = 0.074 \text{ mg cm}^{-2}$) (Fig. 3). In a tick climbing repellency assay using *Rhipicephalus appendiculatus*, the oils showed high repellent activity (chemotype A $RD_{50} = 0.005 \text{ mg}$ and chemotype B $RD_{50} = 0.0012 \text{ mg}$) (Fig. 4).

2.3. Repellent activity of individual compounds and synthetic blends of essential oil components

Repellency testing against *An. gambiae* using purified (4*a*S,7*S*,7*a*R) and (4*a*S,7*S*,7*a*S)-nepetalactone isomers revealed overall lower repellent activity compared to the chemotype A and B oils (Fig. 5). Testing of binary mixtures of the (4*a*S,7*S*,7*a*R) and (4*a*S,7*S*,7*a*S) isomers across a range of ratios, but all at the same overall dose (0.1 mg), revealed not only synergistic activity for mixtures, but also a ratio-dependent effect, with lower activity for pure isomers and equivalent or near-equivalent mixtures (Fig. 6). Furthermore, a binary mixture of (4*a*S,7*S*,7*a*R) and (4*a*S,7*S*,7*a*S) isomers, in a ratio equivalent to that found in chemotype B oil, was less repellent than the oil itself, when tested at two doses equivalent to 0.1 and 0.01 mg chemotype B oil (Fig. 7). Addition of (*E*)-(1*R*,9*S*)-caryo-




Fig. 1. Coupled GC–MS analysis of *Nepeta cataria* chemotype A and B oils obtained by steam distillation of wet-harvested materials. **1** = (4*a*S,7*S*,7*a*R)-nepetalactone; **2** = (4*a*S,7*S*,7*a*S)-nepetalactone; **3** = (*E*)-(1*R*,9*S*)-caryophyllene.




Fig. 2. Mean (\pm SE) percentage protective efficacy provided by *Nepeta cataria* chemotype A and B oils, and *N,N*-diethyltoluamide (DEET) against *Anopheles gambiae* s.s. mosquitoes, using a WHO-approved topical application bioassay.




Fig. 3. Mean (\pm SE) percentage protective efficacy provided by *Nepeta cataria* chemotype A and B oils against *Culex quinquefasciatus* mosquitoes, using a WHO-approved topical application bioassay.




Fig. 4. Mean (\pm SE) percentage repellency provided by *Nepeta cataria* chemotype A and B oils, against *Rhipicephalus appendiculatus* ticks, using a climbing repellency bioassay.

phyllene to the mixture at the level found in chemotype B oil, and testing of the three-component blend, showed that this blend had the same activity as chemotype B oil (Fig. 7).




Fig. 5. Mean (\pm SE) percentage protective efficacy provided by pure (4*aS*,7*S*,7*aR*) and (4*aS*,7*S*,7*aS*)-nepetalactone isomers against *Anopheles gambiae* s.s. mosquitoes, using a WHO-approved topical application bioassay.




Fig. 6. Mean (\pm SE) percentage protective efficacy provided by pure (4*aS*,7*S*,7*aR*) and (4*aS*,7*S*,7*aS*)-nepetalactone isomers and binary mixtures against *Anopheles gambiae* s.s. mosquitoes, using a WHO-approved topical application bioassay. Letters represent different blends of (4*aS*,7*S*,7*aR*) and (4*aS*,7*S*,7*aS*) isomers, respectively: A (1:0), B (4:1), C (3:1), D (3:2), E (5:4), F (1:1), G (4:5), H (2:3), I (1:3), J (1:4) and K (0:1). Means with same letter are not significantly different (Proc GLM, LSD test, $P < 0.05$).

In field trapping assays with *Dermanyssus gallinae*, pre-conditioning of traps with *D. gallinae* significantly increased the level of trap capture (Fig. 8; treatment (i) versus (ii) ($P < 0.05$)). Addition of the chemotype A and B oils, and a combination of the two, to traps pre-conditioned with *D. gallinae*, all resulted in a significant reduction of *D. gallinae* trap capture (Fig. 8; treatment (ii) versus (iii) ($P < 0.05$), (iv) ($P < 0.05$) and (v) ($P < 0.01$)). No significant difference was found between treatments (iii), (iv) and (v).

3. Discussion

The results in this study confirm that the essential oils obtained from two chemotypes of the catmint, *N. cataria* (Birkett and Pickett, 2003), are significantly repellent towards sub-Saharan populations of *An. gambiae* and *Cx. quinquefasciatus* mosquitoes, in line with mosquito repellency studies conducted in other global regions. The data suggest that pure (4*aS*,7*S*,7*aR*) and (4*aS*,7*S*,7*aS*)-nepetalactone have lower activity compared to oils, when tested individually and in a binary mixture at a ratio found in the oil, and that full




Fig. 7. Mean (\pm SE) percentage protective efficacy provided by *Nepeta cataria* chemotype B oil, pure (4aS,7S,7aR) and (4aS,7S,7aS)-nepetalactone and (*E*)-(1*R*,9*S*)-caryophyllene against *Anopheles gambiae* s.s. mosquitoes, using a WHO-approved topical application bioassay. Treatment (i) chemotype B oil (0.1%), (ii) (4aS,7S,7aR) and (4aS,7S,7aS)-nepetalactone at equivalent levels found in chemotype B oil at one dose (0.1 mg dose), (iii) (4aS,7S,7aR), (4aS,7S,7aS)-nepetalactone and (*E*)-(1*R*,9*S*)-caryophyllene at equivalent levels found in chemotype B oil (0.1 mg dose), (iv) chemotype B oil (0.01 mg dose), (v) (4aS,7S,7aR) and (4aS,7S,7aS)-nepetalactone at equivalent levels found in chemotype B oil (0.01 mg dose), (vi) (4aS,7S,7aR), (4aS,7S,7aS)-nepetalactone and (*E*)-(1*R*,9*S*)-caryophyllene at equivalent levels found in chemotype B oil (0.01 mg dose). Means with same letter are not significantly different (Proc GLM, LSD test, $P < 0.05$).




Fig. 8. Mean (\pm SE) percentage trap catches of *Dermanyssus gallinae* mites in a chicken stable ($n = 5$). (i) Clean trap; (ii) mite-conditioned trap; (iii) mite-conditioned trap + *Nepeta cataria* chemotype A oil (5 mg); (iv) mite-conditioned trap + *Nepeta cataria* chemotype B oil (5 mg); (v) mite-conditioned trap + *Nepeta cataria* chemotype A and B oils (5 mg each). Data were analysed by log transformation ($\log(x + 1)$, Zar, 1974) of the means for each treatment, and comparison of treatment (ii) with treatments (i), (iii), (iv) and (v), using a pair-wise Student's *t*-test (* = $P < 0.05$; ** = $P < 0.01$).

repellent activity is observed when the sesquiterpene hydrocarbon (*E*)-(1*R*,9*S*)-caryophyllene is included in a synthetic blend with the nepetalactone isomers. Furthermore, the results in this study show a potent repellent effect of *N. cataria* oils against the brown ear tick, *R. appendiculatus*. To our knowledge, this is the first published report of *N. cataria* repellency against a tick species, thus extending the range of pest arthropods that *N. cataria* can be potentially used against to provide protection.

N. cataria oil has been reported as a potent repellent for the northern house mosquito, *Cx pipiens*, comparing favourably with DEET (Schultz et al., 2006), but studies elsewhere have shown that the purified (4aS,7S,7aR) and (4aS,7S,7aS)-nepetalactones are less effective than DEET (Chauhan et al., 2005). Our data are in line with these previously reported observations. The presence of (*E*)-

(1*R*,9*S*)-caryophyllene in *N. cataria* floral volatiles was reported (Schultz et al., 2004), but no investigation of the role of this component in contributing towards repellency of *N. cataria* oil was reported. Therefore, our study is the first to demonstrate that (*E*)-(1*R*,9*S*)-caryophyllene contributes towards the overall repellency of *N. cataria* oil. Furthermore, the data highlight the fact that mixtures of compounds contribute to greater repellent activity versus *An. gambiae*, as demonstrated previously with essential oils collected from Kenyan plant species (Omolo et al., 2004).

Previous work on the seasonal occurrence of nepetalactone isomer content in *N. cataria* reported variation in the ratio of (4aS,7S,7aR) and (4aS,7S,7aS)-nepetalactone over the course of several weeks, but no clear relationship between ratios and cockroach repellent activity was found (Schultz et al., 2004). The repellency data with oils from *N. cataria* chemotypes A and B in this study reveal differences in repellent activity towards both *An. gambiae* and *Cx. quinquefasciatus*, possibly arising as a consequence of differences in the ratios of nepetalactone isomers between the two oils, as revealed by GC–MS analysis. This hypothesis was supported by the testing of binary mixtures across a range of ratios at a single dose (0.1% mg). Mixtures of (4aS,7S,7aR) and (4aS,7S,7aS)-nepetalactone were consistently more repellent than the isomers when tested alone, apart from equivalent or near-equivalent mixtures, for which repellent activity was surprisingly lower. This suggests that the observed synergistic activity of the nepetalactone isomers is composition dependent.

There is a growing amount of evidence to support the hypothesis that host location by phytophagous insects involves the olfactory perception of ratios of ubiquitous plant compounds (Bruce et al., 2005), as exemplified by the identification of host plant compounds for the black bean aphid, *Aphis fabae* (Webster et al., 2008). It is hypothesised that in conveying a strong plant cue, botanicals interfere with host location by biting and blood-sucking arthropods (Pickett et al., 2010), thus rationalising the development of botanical repellents. The ecological basis for avoidance of *N. cataria*, including specific mixtures of nepetalactone isomers, by mosquitoes is unclear, and further work is required to provide a greater understanding of *N. cataria* and nepetalactone repellency at the neurophysiological and behavioural level.

The use of botanically-derived materials (oils and resins) for the control of African tick pests has been studied elsewhere (Birkett et al., 2008; Mwangi et al., 1995; Ndungu et al., 1995; Lwande et al., 1999). As described above, to our knowledge, this study is the first to report a repellent effect for *N. cataria* against ticks, with activity comparable with other essential oils (Lwande et al., 1999). The gum resin from *Commiphora holtziana* (Burseraceae), known as gum haggard, has been shown to be repellent for the red poultry mite, *D. gallinae*, which is another major acarine pest, affecting chicken production systems (Birkett et al., 2008). The ability of (4aS,7S,7aR) and (4aS,7S,7aS)-nepetalactone to interfere with *D. gallinae* behaviour in this study demonstrates the potential for using *N. cataria* oils and compounds as general purpose acarine repellents, especially in organic production systems, where the use of broad-spectrum toxicants, i.e. acaricides, is prohibited.

In summary, the results of this study reinforce the potential for using *N. cataria*-derived materials in protection against Afro-tropical mosquito pathogen vectors, but also demonstrate their potential to be used to protect against ixodid ticks, and other acarine pests. Furthermore, *Nepeta* spp. essential oils appear to offer greater protection compared to pure nepetalactone isomers, due to the presence of minor components, e.g. (*E*)-(1*R*,9*S*)-caryophyllene, that contribute to their overall activity. However, it should be noted that variation in the activity of oils from different chemotypes suggests that the screening and selection of specific varieties is crucial for developing product(s) with optimal activity.

4. Experimental

4.1. *Nepeta cataria* extraction and chemicals

The essential oils from flowering harvested *N. cataria* (chemotypes A and B) were prepared by steam distillation using cyclohexane as the co-solvent as described previously (Botanix Ltd., Paddock Wood, Kent) (Birkett and Pickett, 2003). Authentic samples of (4aS,7S,7aR) and (4aS,7S,7aS)-nepetalactone (1.5 g; >99% purity by GC) were obtained by liquid chromatographic isolation from *N. cataria* A and B oils over silica (40–60 mesh, Sigma–Aldrich, Gillingham, UK), using hexane and diethyl ether (1:9) as eluant, and their identities confirmed by comparison of ¹H and ¹³C NMR data with literature values (Dawson et al., 1996). (*E*)-(1*R*,9*S*)-Caryophyllene (>99% by GC) was purchased from Sigma–Aldrich (Gillingham, UK).

4.2. GC–MS analysis

GC–MS analysis of oils collected from *N. cataria* chemotypes A and B was performed using a fused silica capillary column (50 m × 0.32 mm i.d., film thickness 0.52 μm, DB-1, J&W Scientific), attached to a cool on-column injector, which was directly coupled to a magnetic sector mass spectrometer (Autospec Ultima, Fisons Instruments, Manchester, UK). Ionization was by electron impact (70 eV, source temperature 250 °C). Helium was the carrier gas. The oven temperature was maintained at 30 °C for 5 min, and then programmed at 5 °C/min to 250 °C. Tentative identifications were made by comparison of spectra with mass spectral databases (NIST, 2005), and confirmed by peak enhancement on GC using authentic compounds (Pickett, 1990).

4.3. Mosquitoes, ticks and mites

Mosquitoes used in this study were obtained from laboratory colonies reared at ICIPE Duduville campus, Nairobi, Kenya, according to a WHO protocol (1996). *Anopheles gambiae* s.s. (Ifakara stain) was cultured in 1998 from specimens originally obtained from Njage, 70 km from Ifakara, south-east from Tanzania. A colony of *Culex quinquefasciatus* (Nairobi strain) was established in 2006 from field collections of larvae from typical culicine pools. Larvae were reared in plastic trays (39 × 28 × 14 cm) at a density of ~500 larvae in 3 l of distilled water at 32–36 °C and fed daily with TetraMin®. The adults were kept in cubic cages (30 × 30 × 30 cm) in separate rooms maintained at 26–28 °C, 70–80% humidity and photoperiod of 12:12 (L:H), and were provided with 6% glucose solution to feed on ad libitum. The females were fed on human blood from a volunteer's forearm thrice a week. Approval for feeding mosquitoes on human volunteers was obtained from the Kenya National Ethical Review Board (protocol number KEMRI/RES/7/3/1). Adult ticks of *Rhipicephalus appendiculatus* were obtained from a colony established at ICIPE, Nairobi, since 2002, which had originated from a culture obtained from International Livestock Research Institute (ILRI), Nairobi, Kenya. The ticks were reared under conditions and management initially developed by Bailey (1960) and then optimised by Irvine and Brocklesby (1970). Red poultry mites, *Dermanyssus gallinae*, were collected from poultry sheds at Lovsta Research Station, Swedish University of Agricultural Sciences, Uppsala, using cardboard traps (Nordenfors and Chirico, 2001), and used directly for field experiments.

4.4. Mosquito repellency assay

Assays were performed with 5–7 day old female *An. gambiae* or *Cx. quinquefasciatus* that had been starved for 18 h, but previ-

ously fed on 6% glucose solution. Six human volunteers were selected from those who showed mild or no allergic reaction to mosquito bites or oils/chemicals. They had no contact with lotions, perfumes, oils or perfumed soaps on the day of the experiment. Participation of human volunteers in the repellency assays was part of the approval obtained from the Kenya National Ethical Review Board, protocol number KEMRI/RES/7/3/1. Assays were carried out in aluminium-frame cages (50 × 50 × 50 cm) with cotton stockinet sleeves on the front (WHO, 1996) in a 7 × 5 × 3 m room at 30–32 °C and relative humidity of 65–80%, with 50 female *An. gambiae* or *Cx. quinquefasciatus* starved for 18 h in each cage. Test solutions (0.5 ml) were dispensed on one of the forearms of a volunteer from the wrist to the elbow. The rest of the hand was covered with a glove. Acetone (0.5 ml, HPLC grade) was dispensed on the other forearm to serve as a control. The control and treated arms were interchanged regularly to eliminate bias. The control arm was introduced into the cage 3 min after releasing mosquitoes (to allow the mosquitoes to settle) and kept there for 3 min. The number of insects that landed on that arm during the test duration was recorded. The treated arm was then introduced into the cage for the same period of time and the number of landing insects recorded. The different concentrations of each test sample were tested sequentially starting with the lowest dose. Repellency data from six replicates, expressed as protective efficacy (PE) at each dose, were calculated using the formula, PE = (% control mean – % test mean) / % control mean. The data were transformed and subjected to ANOVA. Means were ranked using the Student–Newman–Keuls (SNK) test.

The following materials were tested: (i) *N. cataria* chemotype A and B oils, between 0.01 and 10 mg versus *An. gambiae*, and 0.001 and 10 mg versus *Cx. quinquefasciatus*; (ii) (4aS,7S,7aR) and (4aS,7S,7aS)-nepetalactone, tested individually between 0.01 and 1 mg versus *An. gambiae*; (iii) binary mixtures of (4aS,7S,7aR)-nepetalactone + (4aS,7S,7aS)-nepetalactone in the following ratios: 1:0, 4:1, 3:1, 3:2, 5:4, 1:1, 4:5, 2:3, 1:3, 1:4 and 0:1, all at a 0.1 mg dose versus *An. gambiae*; (iv) *N. cataria* chemotype B oil at 0.1 and 0.01 mg, (4aS,7S,7aR)-nepetalactone + (4aS,7S,7aS)-nepetalactone in a ratio and dose equivalent to chemotype B oil (at 0.1 and 0.01 mg) and (4aS,7S,7aR)-nepetalactone + (4aS,7S,7aS)-nepetalactone + (*E*)-(1*R*,9*S*)-caryophyllene in a ratio and dose equivalent to chemotype B oil (at 0.1 and 0.01 mg). RD₅₀ values for the chemotype A and B oils were obtained by probit analysis using the PE values obtained from replicated experiments (Omolo et al., 2004).

4.5. Tick climbing assay

A dual choice tick repellency climbing assay (Wanzala et al., 2004) that exploits the behaviour of *R. appendiculatus* to climb up grass stems to await potential passing hosts (Browning, 1976; Chiera, 1985) was used. *N. cataria* chemotype A and B oils were diluted serially from 1000 parts per million (ppm) to 0.05 ppm using methylene chloride (GC grade). An aliquot of each dose (20 μl) was applied onto a filter paper strip, with an equivalent volume of methylene chloride being added to control filter paper strips. Five mixed adult *R. appendiculatus* were introduced at the base of a climbing assay set-up (Wanzala et al., 2004) and their movements observed. Observations were made over a 1 h period at 15, 30, 45 and 60 min. Twelve replicates for each dose were carried out. The repellency was calculated using the formula: (number of ticks in control arm – number of ticks in treated arm / total responding ticks) × 100. Dose-response data were subjected to probit analysis using the % repellency values obtained from replicated experiments (Wanzala et al., 2004).

4.6. Chicken mite field experiment

Cardboard traps (Nordenfors and Chirico, 2001) were used to assess the impact of different treatments on catches of red poultry mites, *D. gallinae*, in a chicken stable (Fig. 8). Five treatments (i)–(v) were assessed. Treatment (i) comprised clean traps, which were used as a passive control, whereas treatment (ii) comprised traps which had been subject to mite conditioning by placing them overnight together with 500 freshly caught mites. The conditioning reinforces the mite arrestment behaviour and significantly increases trap catches (Fig. 8). Treatments (iii) and (iv) comprised mite-conditioned traps with *N. cataria* chemotype A and B oils added, respectively (5 mg per trap). Treatment (v) comprised a mite-conditioned trap with both oils added (5 mg of each). The traps were randomly distributed in the chicken stable, left for a period of 24 h, then collected and placed in a freezer to permit a convenient counting of individuals caught. Data were analysed by log transformation ($\log(x + 1)$, Zar, 1974) of the means for each treatment, and comparison of treatment (ii) with treatments (i), (iii), (iv) and (v) using a pair-wise Student's-*t*-test. The experiment was replicated five times.

Acknowledgements

This work was funded under the BBSRC LINK Competitive Industrial Materials from Non-Food Crops (CIMNFC) programme. Rothamsted Research receives grant-aided support from the Biotechnology and Biological Sciences Research Council (BBSRC) of the United Kingdom. Dr. J. Chirico, Swedish National Veterinary Institute (SVA), is greatly acknowledged for providing support with facilities for the red poultry mite experiment.

References

- Amer, A., Mehlhorn, H., 2006. Repellency effect of forty-one essential oils against *Aedes*, *Anopheles* and *Culex* mosquitoes. *Parasitology Research* 99, 478–490.
- Bailey, K.P., 1960. Notes on the rearing of *Rhipicephalus appendiculatus* and their infection with *Theileria parva* for experimental transmission. *Bulletin of Epizootic Diseases in Africa* 8, 33–43.
- Bernier, U.R., Furman, D.L., Kline, D.L., Allan, S.A., Barnard, D.R., 2005. Comparison of contact and spatial repellency of catnip oil and *N,N*-diethyl-3-methylbenzamide (DEET) against mosquitoes. *Journal of Medical Entomology* 42, 306–311.
- Birkett, M.A., Pickett, J.A., 2003. Aphid sex pheromones: from discovery to commercial production. *Phytochemistry* 62, 651–656.
- Birkett, M.A., Al Abassi, S., Kröber, T., Chamberlain, K., Hooper, A.M., Guerin, P.A., Pettersson, J., Pickett, J.A., Slade, R., Wadhams, L.J., 2008. Antiectoparasitic activity of the gum resin, gum haggard, from the East African plant, *Commiphora holtziana*. *Phytochemistry* 69, 1710–1715.
- Browning, T.O., 1976. The aggregation of questing ticks, *Rhipicephalus pulchellus* on grass stems, with observations on *Rhipicephalus appendiculatus*. *Physiological Entomology* 1, 107–114.
- Bruce, T.J.A., Wadhams, L.J., Woodcock, C.M., 2005. Insect host location: a volatile situation. *Trends in Plant Science* 10, 269–274.
- Cantrell, C.L., Klun, J.A., Bryson, C.T., Kobaisy, M., Duke, S.O., 2005. Isolation and identification of mosquito bite deterrent terpenoids from leaves of American (*Callicarpa americana*) and Japanese (*Callicarpa japonica*) Beautyberry. *Journal of Agricultural and Food Chemistry* 53, 5948–5953.
- Carroll, J.F., Cantrell, C.L., Klun, J.A., Kramer, M., 2007. Repellency of two terpenoid compounds isolated from *Callicarpa Americana* (Lamiaceae) against *Ixodes scapularis* and *Amblyomma americanum* ticks. *Experimental and Applied Acarology* 41, 215–224.
- Chauhan, K.R., Klun, J.A., Debboun, M., Kramer, M., 2005. Feeding deterrents of catnip oil components compared with two synthetic amides against *Aedes aegypti*. *Journal of Medical Entomology* 42, 643–646.
- Chiera, W.J., 1985. The distribution and climbing behaviour of *Rhipicephalus appendiculatus* Neumann on grass stems. *Insect Science and its Application* 6, 213–215.
- Chou, J.T., Rossignol, P.A., Ayres, J.W., 1997. Evaluation of commercial insect repellents on human skin against *Aedes aegypti* (Diptera: Culicidae). *Journal of Medical Entomology* 34, 624–630.
- Dawson, G.W., Pickett, J.A., Smiley, D.W.M., 1996. The aphid sex pheromone cyclopentanoids: synthesis in the elucidation of structure and biosynthetic pathways. *Bioorganic and Medicinal Chemistry* 4, 351–361.
- Debboun, M., Frances, S.P., Strickman, D.A. (Eds.), 2006. *Insect Repellents: Principles, Methods and Uses*. CRC Press, Florida, USA.
- Food and Agriculture Organisation of the United Nations, 2005. Impact of Climate Change, Pests and Diseases on Food Security and Poverty Reduction. 31st Session of the Committee of World Food Security, 23–26 May.
- Irvine, A.D., Brocklesby, D.W., 1970. Rearing and maintaining *Rhipicephalus appendiculatus* in the laboratory. *Journal of Institute for Animal Technology* 21, 106–112.
- Lindsay, L.R., Surgeoner, G.A., Heal, J.D., Gallivan, G.J., 1996. Evaluation of the efficacy of 3% citronella candles and 5% citronella incense for protection against field populations of *Aedes* mosquitoes. *Journal of the American Mosquito Control Association* 12, 293–294.
- Logan, J.G., Birkett, M.A., 2007. Semiochemicals for biting fly control: their identification and exploitation. *Pest Management Science* 63, 647–657.
- Lwande, W., Ndakala, A.J., Hassanali, A., Moreka, L., Nyandat, E., Ndungu, M., Amiani, H., Gitu, P.M., Malonza, M.M., Punyua, D.K., 1999. *Gynandropsis gynandra* essential oil and its constituents as tick (*Rhipicephalus appendiculatus*) repellents. *Phytochemistry* 50, 401–405.
- Mwangi, E.N., Hassanali, A., Essuman, S., Myandat, E., Moreka, L., Kimondo, M., 1995. Repellent and acaricidal properties of *Ocimum suave* against *Rhipicephalus appendiculatus* ticks. *Experimental and Applied Acarology* 19, 11–18.
- Ndungu, M., Lwande, W., Hassanali, A., Moreka, L., Chabira, S.C., 1995. *Cleome monophylla* essential oil and its constituents as tick (*Rhipicephalus appendiculatus*) and maize weevil (*Sitophilus zeamais*) repellents. *Entomologia Experimentalis et Applicata* 76, 217–222.
- Nishimura, H., Satoh, A., 1999. Potent mosquito repellents from the leaves of *Eucalyptus* and *Vitex* plants. In: Cutler, H.G., Cutler, S.J. (Eds.), *Biologically Active Natural Products: Agrochemicals*. CRC Press, pp. 137–146.
- NIST, 2005. NIST/EPA/NIH Mass Spectral Library Version 2.0. Office of the Standard Reference Data Base, National Institute of Standards and Technology, Gaithersburg, Maryland.
- Nordenfors, H., Chirico, J., 2001. Evaluation of a sampling trap for *Dermanyssus gallinae* (Acari: Dermanyssidae). *Journal of Economic Entomology* 94, 1617–1621.
- Odalo, J.O., Omolo, M.O., Malebo, H., Angira, J., Njeru, P.M., Ndiege, I.O., Hassanali, A., 2005. Repellency of essential oils of some plants from the Kenyan coast against *Anopheles gambiae*. *Acta Tropica* 95, 210–218.
- Omolo, M., Okinyo, D., Ndiege, I.O., Lwande, W., Hassanali, A., 2004. Repellency of essential oils of some Kenyan plants against *Anopheles gambiae*. *Phytochemistry* 65, 2797–2802.
- Peterson, C., Coats, J., 2001. Insect repellents – past, present and future. *Pesticide Outlook* 12, 154–158.
- Pickett, J.A., 1990. Gas chromatography–mass spectrometry in insect pheromone identification: three extreme cases. In: McCaffery, A.R., Wilson, I.D. (Eds.), *Chromatography and Isolation of Insect Pheromones and Hormones*. Plenum Press, New York, pp. 209–309.
- Pickett, J.A., Birkett, M.A., Dewhurst, S.V., Logan, J.G., Omollo, M.O., Torto, B., Pelletier, J., Syed, Z., Leal, W.S., 2010. Chemical ecology of animal and human pathogen vectors in a changing global climate. *Journal of Chemical Ecology* 36, 113–121.
- Polsomboon, S., Grieco, J.P., Achee, N.L., Chauhan, K.R., Tanasinchayakul, S., Pothikasikorn, J., Chareonviriyaphap, T., 2008. Behavioural responses of catnip (*Nepeta cataria*) by two species of mosquitoes, *Aedes aegypti* and *Anopheles harrisoni*, in Thailand. *Journal of the American Mosquito Control Association* 24, 513–519.
- Schultz, G., Simbro, E., Belden, J., Zhu, J., Coats, J., 2004. Catnip, *Nepeta cataria* (Lamiaceae) – a closer look: seasonal occurrence of nepetalactone isomers and comparative repellency of three terpenoids to insects. *Environmental Entomology* 33, 1562–1569.
- Schultz, G., Peterson, C., Coats, J., 2006. Natural insect repellents: activity against mosquitoes and cockroaches. *ACS Symposium Series* 927, 168–181.
- Spero, N.C., Gonzalez, Y.I., Scialdone, M.A., Hallahan, D., 2008. Repellency of hydrogenated catmint oil formulations to black flies and mosquitoes in the field. *Journal of Medical Entomology* 45, 1080–1086.
- Wanzala, W., Sika, N.F.K., Gule, S., Hassanali, A., 2004. Attractive and repellent host odours guide ticks to their respective feeding sites. *Chemoecology* 14, 229–232.
- Webb, C.E., Russell, R.C., 2007. Is the extract from the plant catmint (*Nepeta cataria*) repellent to mosquitoes in Australia? *Journal of the American Mosquito Control Association* 23, 351–354.
- Webster, B., Bruce, T., Dufour, S., Birkemeyer, C., Birkett, M., Hardie, J., Pickett, J., 2008. Identification of volatile compounds used in host location by the black bean aphid, *Aphis fabae*. *Journal of Chemical Ecology* 34, 1153–1161.
- WHO, 1996. Report of the WHO Informal Consultation on the Evaluation and Testing of Insecticides. World Health Organisation, Geneva, pp. 32–36, 5–52.
- WHO, 2000. The World Health Report 2000. Health Systems: Improving Performance. World Health Organisation, Geneva, <www.who.int/whr/2000/en/statistics.htm>.
- Zar, J.H., 1974. *Biostatistical Analysis*. Prentice-Hall, London, p. 184.
- Zhang, A., Klun, J., Wang, S., Carroll, J.F., Debboun, M., 2009. Isolongifolenone: a novel sesquiterpene repellent of ticks and mosquitoes. *Journal of Medical Entomology* 46, 100–106.